

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)			2. REPORT TYPE Technical Papers		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)					5d. PROJECT NUMBER 2303	
					5e. TASK NUMBER M2C8	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048					8. PERFORMING ORGANIZATION REPORT	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT A	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Leilani Richardson	
					19b. TELEPHONE NUMBER (include area code) (661) 275-5015	
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified				

C P

TP-1998-096

P98-096

MEMORANDUM FOR IN-HOUSE PUBLICATIONS

FROM: PROI (TI) (STINFO)

30 Apr 98

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-1998-096

J. Harper, J. Sheehy, and Bill Larson "Isolation of Carbon Atoms in Cryogenic Solids"

HEDM Conference Presentation

(Statement A)

Isolation of Carbon Atoms in Cryogenic Solids

J. Harper, J. A. Sheehy, and C. W. Larson

Propulsion Sciences and Advanced Concepts Division
Air Force Research Laboratory
Propulsion Directorate
Edwards AFB, CA 93524-7680

20021122 005

AFOSR HIEDM Contractor's Conference
Monterey, CA 20-22 May 1998


DISTRIBUTION STATEMENT A:
Approved for Public Release
Distribution Unlimited

Project Goals

- Increase production of carbon atoms from furnace source
- Prevent condensation of carbon atoms in matrix
- Develop general approach to make other HEDM propellants
- Identify new molecules through spectroscopy
- Verify IR assignments to aid in interpretation of results

Progress toward 5% carbon atoms in solid H₂
(To yield 469s I_{sp} propellant, compared to 389s for LOX/LH₂)


Experiment



- Higher oven temperature increases carbon atom production
- Smaller iris in heat shield reduces radiative heat load on matrix from oven
- Slower argon flow rate (to a limit) increases concentration of carbon atoms
- Contaminates scavenge carbon atoms; high vacuum (10^{-8} Torr) critical

FTIR Spectroscopy

- Method for direct detection of carbon atoms unavailable
- Indirect atom detection by analysis of IR spectra taken before and after annealing of the carbon/argon matrix
 - Absolute column densities (molecules cm⁻²) are determined for each carbon cluster from Beer's Law.
 - Identify and track all important species that grow or disappear during annealing processes.
 - From successive cluster distributions one can calculate the quantity of carbon species that were not detected in the as-deposited matrix IR spectrum. This increase in the total equivalent carbon atom density is attributed to carbon atoms that are “invisible” to IR and were thus not accounted for in the original IR spectrum.



Ja218d62.axg Apr. 28, 1998 1:52:15 PM

0.5-

0.4-

0.3-

0.2-

0.1-

0.0-

1997.7

1952.0

1912.9

1893.9

1849.8

1803.3

1745.9

1714.1

1699.7 - 1694.5

1623.5

1544.9

1600


1700


1800


1900

2000


2100







Ja315d4.aug Feb. 26, 1998 5:04:46 PM




Computational Support

- The vibrational frequencies, infrared signal intensities, absorption cross section, and isotopic frequency shifts for carbon clusters (C_n , $n < 13$) are determined using B3LYP and CCSD(T) with cc-pVQZ and higher basis sets [6-31G*, 6-311G, 6-311G(2d), 6-311G(2df)].
- These results aid greatly in the interpretation of infrared spectra of the matrices. Cyclic C_6 , cyclic C_8 , and tentatively cyclic C_{10} molecules have been identified through theoretical and experimental validation. IR frequency shifts due to matrix effects have been identified for other molecules, such as C_{13} , which are well characterized in the gas phase.

Cyclic C_8

B3LYP/cc-pVDZ optimized structure (C_{4h} symmetry)



Future Work

- Switch to liquid helium cryostat to perform experiments at 4K . Using argon one can measure the accretion time at this temperature to compare with that from the experiments performed at 10K , before moving on to depositions in H_2 . A limitation to the liq. He cryostat is that the matrix in the system cannot be easily annealed.
- If results look promising, it will be worthwhile to invest in a method of direct detection of carbon atoms to use in the 4K system.

Conclusions

- Highest equivalent carbon atom concentration attained is 0.1 mol%.
- Hotter oven produces more carbon atoms
- FTIR is an effective way to derive the concentration of carbon atom equivalents in the matrices
- The temperature of the substrate is critical in preventing condensation of carbon atoms
- Unknown peaks in IR spectra have been assigned:
 - weaker vibrational mode of C₁₃, cyclic C₈, tentatively cyclic C₁₀
- Cyclic species dominate high density matrices